


Evito® Fungicide

With Xylem Pro Technology™


Peanut - Yield (2008)

2008 Large Plot Trials - Both strobis applied at 60 & 90 days




Peanut - Rhizoctonia (2008)

2008 Large Plot Trials - Both strobis applied at 60 & 90 days



Peanut - White Mold (2008)

2008 Large Plot Trials - Both strobis applied at 60 & 90 days



Active ingredient – fluoxastrobin

Strobilurin fungicide

Formulation – 480 SC (4 lb ai/gal)

Attributes

- Broad-spectrum
- Strobilurin-induced yield increases
- Xylem mobile (moves with water flow)
- Better rainfastness for leaf spot
- Excellent soil-borne disease activity
- Excellent crop safety

Evito[®] Fungicide

With Xylem Pro Technology™

Application Rates and Timing for EVITO[®] Fungicide

Diseases	Use Rate	Application Guidelines
Early leaf spot <i>(Cercospora arachidicola)</i> Late leaf spot <i>(Cercosporidium personatum)</i> Leaf rust <i>(Puccinia arachidis)</i>	3.8 to 5.7 fl oz/Acre (0.12 to 0.18 lbs ai/Acre)	For optimum results, begin application preventively. Apply as needed on a 14-day interval. To limit the potential for development of disease resistance: <ul style="list-style-type: none"> • In areas with typically 1-4 sprays per year, alternate every application of a solo QoI fungicide with at least one application of another effective mode of action fungicide. • In areas with typically 5 or more fungicide sprays per year, a maximum of 2 sequential applications of a QoI fungicide followed by at least an equal number of another effective mode of action fungicide.
Disease Suppression		Improved white mold control can be obtained with higher rates of EVITO (5.7 fl oz/Acre, per federal label) or with combinations (tank-mixed or alternating applications) with other products labels for white mold.
Stem rot		When targeting white mold at the lower rate of 3.8 fl oz/Acre, we suggest a tank-mix with tebuconazole for best results.
White mold		
Southern blight <i>(Sclerotium rolfsii)</i>		
Rhizoctonia limb rot <i>(Rhizoctonia solani)</i>		


It pays to protect with EVITO Fungicide.

Performance. Consistency. Flexibility. The benefits of using EVITO on peanuts pay off where it matters most — at harvest.

- Broad-spectrum control of early and late leaf spot, leaf rust, white mold and *Rhizoctonia* limb rot
- Fast-acting, preventive control
- Short Restricted Entry Interval (REI) of just 12 hours
- 14-day Pre-harvest Interval (PHI)
- EVITO 4lb active ingredient per gallon vs. 2.08 / 2.09 active ingredient per gallon competitors

Rainfastness of Fluoxastrobin (2008)

8 greenhouse trials



Always read and follow label directions. EVITO and the EVITO logo are registered trademarks of Arysta LifeScience North America, LLC. The "Xylem Pro Technology" slogan is a trademark of Arysta LifeScience North America, LLC. Arysta LifeScience and the Arysta LifeScience logo are registered trademarks of Arysta LifeScience Corporation. ©2009 Arysta LifeScience North America, LLC. EVT-053