

Kanemite[®] Miticide

Might Versus Mites[™]



For fast-acting and long-lasting control of citrus red mite, Texas citrus mite and other damaging mites.

University trials and actual grower use prove that nothing controls citrus red mite (*Panonychus citri*), Texas citrus mite (*Eutetranychus banksi*) and other damaging mites like KANEMITE[®] 15 SC Miticide from Arysta LifeScience North America. That's because KANEMITE destroys damaging mites at every life stage, from eggs through adults.

In addition, KANEMITE provides outstanding residual control of damaging mites while preserving predacious mites and other beneficial species. That's important, since predators and parasites are vital in developing a successful, season-long mite control program.

Finally, KANEMITE Miticide also features the unique active ingredient acequinocyl, with no known cross-resistance to other commonly used miticides. This fact makes KANEMITE the ideal choice for managing the growing threat of mite resistance.

When you add it all up, it's easy to see why KANEMITE is such an exceptional value. The kind of value you can count on for higher yields of more marketable fruit.

New EPA Approved
Expanded Label

Kanemite[®] Miticide

Might Versus Mites[™]

Application Rates and Timing for Citrus

Pest		Rate of Product per Acre	Application Timing
Twospotted spider mite (<i>Tetranychus urticae</i>)	Pacific spider mite (<i>Tetranychus pacificus</i>)	21.0-31.0 fl oz/A (0.2-0.3 lb ai/acre)	To provide maximum control, apply KANEMITE 15 SC before mite populations exceed economic thresholds. Heavy infestations require thorough coverage.
Spruce mite (<i>Oligonychus ununguis</i>)	Texas citrus mite (<i>Eutetranychus banksi</i>)		
European red mite (<i>Panonychus ulmi</i>)	Strawberry spider mite (<i>Tetranychus turkestanii</i>)		
Citrus red mite (<i>Panonychus citri</i>)			

Application Guidelines for Citrus

- Maximum seasonal use rate: Do not apply more than a total of 62 fl oz of KANEMITE 15 SC per acre per season.
- Do not make more than two applications of KANEMITE per season.
- Allow a minimum of 21 days between applications of KANEMITE.
- Use KANEMITE in rotation with other miticide treatments having different modes of action.
- Do not use less than 100 gallons of water volume per acre.
- Do not apply KANEMITE by air.
- Do not apply this product through any type of irrigation system.
- Do not apply within 7 days of harvest.

There's a lot to like about KANEMITE.

Along with effective and selective control of citrus red mite, Texas citrus mite and other damaging mites, KANEMITE Miticide offers you these other value-added benefits:

- Rapid knockdown to stop mite flare-ups before they get out of hand
- Long-lasting residual activity
- A unique mode of action with no known cross-resistance
- Activity on mites at all life stages (eggs, larvae and adults)
- Soft on predacious mites, honeybees and other beneficial species
- A 7-day Pre-harvest Interval (PHI) for application flexibility
- A short 12-hour Restricted Entry Interval (REI)
- A Reduced Risk classification for added peace of mind
- An excellent fit for IPM and resistance management programs
- A convenient flowable formulation for easier handling and mixing

To learn how KANEMITE 15 SC Miticide can help improve your mite control program, contact your crop consultant or retailer today. And for more information about KANEMITE, log on to www.arystalifescience.us/kanemite or call 1-866-761-9397 toll free.

Best Use Guidelines for KANEMITE Miticide

In order to maximize the efficacy of KANEMITE, follow these guidelines:

- Once product is obtained from distributor or warehouse, place in cool dark storage area.
- Do not expose product to direct sunlight or to temperatures above 104 degrees F (40 degrees C).
- When mixing, loading or spraying, minimize product's exposure to sunlight and temperatures above 104 degrees F.
- Apply KANEMITE at 21-31 fl oz of product per acre.
- Apply KANEMITE in a minimum of 100 gallons for sufficient coverage.
- Use of a non-ionic surfactant enhances coverage.
- Reapply KANEMITE according to the above recommendations should mite pressures rebuild, but not before 21 days (since previous application) and no more than two applications per season.

Always read and follow label directions. KANEMITE and the KANEMITE logo are registered trademarks of Arysta LifeScience North America LLC. The "Might Versus Mites" slogan is a trademark of Arysta LifeScience North America LLC. Arysta LifeScience and the Arysta LifeScience logo are registered trademarks of Arysta LifeScience Corporation. ©2008 Arysta LifeScience North America LLC. KAN-076